

MEK2 Rabbit Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP11273

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

 Immunogen
 Recombinant Protein of MEK2

Mol wt N/A

Species reactivity Human

Clonality Polyclonal

Recommended application WB, IHC-p, IF

Concentration 1 mg/ml

Full name Dual specificity mitogen-activated protein kinase kinase 2

Synonyms Dual specificity mitogen-activated protein kinase 2 ;MAP kinase kinase 2 ;MAPKK 2;EC

2.7.12.2; ERK activator kinase 2; MAPK/ERK kinase 2; MEK 2; Dual specificity mitogen-activated protein kinase kinase 2; MAP kinase kinase 2; MAPKK 2; EC 2.7.12.2; ERK activator kinase 2; MAPK/ERK

kinase 2; MEK 2

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.

Recommended Dilution


WB: 1: 500 - 1: 2000 IHC-p: 1: 50 - 1: 300 ELISA: 1: 20000

Not yet tested in other applications.

Images



 $Immun ohistochemical \ analysis \ of paraffin-embedded \ Human \ Colon \ Carcinoma \ Tissue \ using \ MEK2$ Rabbit pAb diluted at 1:200



Western blot analysis of 1) Hela Cell Lysate, 2)293t Cell Lysate using MEK2 Rabbit pAb diluted at 1:2000.

Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com