

HDAC1 Monoclonal Antibody

Description

Product type	Antibody
Code	BT-MCA2452
Host	Mouse
Isotype	Mouse IgG1
Size	100μL, 50μL
Immunogen	Purified recombinant fragment of human HDAC1 (AA: 321-482) expressed in E. Coli.
Mol wt	55.1kDa
Species reactivity	Others
Clonality	Monoclonal
Recommended application	IHC,FCM
Concentration	N/A
Full name	N/A
Synonyms	HD1;RPD3;KDAC1;GON-10;RPD3L1

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis.

Recommended Dilution

WB: 1:500 - 1:2000


IHC-p: 1:200 - 1:1000

FCM: 1:200 - 1:400


ELISA: 1:10000

Not yet tested in other applications.


Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using HDAC1 mAb against human HDAC1 (AA: 321-482) recombinant protein. (Expected MW is 44.6 kDa)




Western blot analysis using HDAC1 mAb against HEK293-6e (1) and HDAC1 (AA: 1-482)-hIgGfc transfected HEK293-6e (2) cell lysate.




Western blot analysis using HDAC1 mouse mAb against NIH/3T3 (1), HeLa (2), Raw264.7 (3), K562 (4), Jurkat (5), C6 (6), and Raji (7) cell lysate.



Flow cytometric analysis of Raji cells using HDAC1 mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human brain tissues using HDAC1 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded bladder cancer tissues using HDAC1 mouse mAb with DAB staining.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com