

CD322 Monoclonal Antibody

Description

Product type Antibody

Code BT-MCA3131

Host Mouse

 $\begin{tabular}{ll} \textbf{Isotype} & Mouse IgG1 \\ \\ \textbf{Size} & 100 \mu L, 50 \mu L \\ \end{tabular}$

Immunogen Purified recombinant fragment of human CD322 (AA: extra 29-238) expressed in E. Coli.

Mol wt 33.2kDa

Species reactivity Human

Clonality Monoclonal

Recommended application WB,FCM

Synonyms JAM2;JAMB;JAM-B;VEJAM;PRO245;VE-JAM;C21orf43

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


This gene belongs to the immunoglobulin superfamily, and the junctional adhesion molecule (JAM) family. The protein encoded by this gene is a type I membrane protein that is localized at the tight junctions of both epithelial and endothelial cells. It acts as an adhesive ligand for interacting with a variety of immune cell types, and may play a role in lymphocyte homing to secondary lymphoid organs. Alternatively spliced transcript variants have been found for this gene.

Recommended Dilution

WB: 1:500 - 1:2000 FCM: 1:200 - 1:400 ELISA: 1:10000


Not yet tested in other applications.

Images




Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)

Western blot analysis using CD322 mAb against human CD322 (AA: extra 29-238) recombinant protein. (Expected MW is 53.3 kDa)



Western blot analysis using CD322 mAb against HEK293 (1) and CD322 (AA: extra 29-238)-hIgGFc transfected HEK293 (2) cell lysate.

Western blot analysis using CD322 mouse mAb against Ramos (1) and HepG2 (2) cell lysate.



Flow cytometric analysis of HL-60 cells using CD322 mouse mAb (green) and negative control (red).

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com