

BCL2 Monoclonal Antibody

Description

Product type Antibody

Code BT-MCA3509

Host Mouse

 Isotype
 Mouse IgG1

 Size
 100μL, 50μL

Immunogen Purified recombinant fragment of human BCL2 (AA: 1-239) expressed in E. Coli.

Mol wt 26.3kDa

Species reactivity Others

Clonality Monoclonal

Recommended application FCM

Concentration N/A
Full name N/A

Synonyms Bc1-2;PPP1R50

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Alternative splicing results in multiple transcript variants.

Recommended Dilution

WB: 1:500 - 1:2000 FCM: 1:200 - 1:400 ELISA: 1:10000


Not yet tested in other applications.

Images




Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)

Western blot analysis using BCL2 mAb against human BCL2 (AA: 1-239) recombinant protein. (Expected MW is $29.2~\mathrm{kDa}$)



Western blot analysis using BCL2 mAb against HEK293-6e (1) and BCL2 (AA: 1-239)-hIgGFc transfected HEK293-6e (2) cell lysate.



Flow cytometric analysis of THP-1 cells using BCL2 mouse mAb (green) and negative control (red).

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com