

WT1 Monoclonal Antibody

Description

| | |
|--------------------------------|--|
| Product type | Antibody |
| Code | BT-MCA3653 |
| Host | Mouse |
| Isotype | Mouse IgG1 |
| Size | 100µL, 50µL |
| Immunogen | Purified recombinant fragment of human WT1 (AA: 1-181) expressed in E. Coli. |
| Mol wt | 49.2kDa |
| Species reactivity | Others |
| Clonality | Monoclonal |
| Recommended application | WB,FCM |
| Concentration | N/A |
| Full name | N/A |
| Synonyms | GUD;AWT1;WAGR;WT33;NPHS4;WIT-2 |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

This gene encodes a transcription factor that contains four zinc-finger motifs at the C-terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with Wilms tumor. This gene exhibits complex tissue-specific and polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal and paternal alleles in different tissues. Multiple transcript variants have been described. In several variants, there is evidence for the use of a non-AUG (CUG) translation initiation codon upstream of, and in-frame with the first AUG. Authors of PMID:7926762 also provide evidence that WT1 mRNA undergoes RNA editing in human and rat, and that this process is tissue-restricted and developmentally regulated.

Recommended Dilution


WB: 1:500 - 1:2000

FCM: 1:200-1:400


ELISA: 1:10000

Not yet tested in other applications.


Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Western blot analysis using WT1 mAb against human WT1 (AA: 1-181) recombinant protein. (Expected MW is 21.2 kDa)



Western blot analysis using WT1 mAb against HEK293 (1) and WT1 (AA: 1-181)-hIgGFc transfected HEK293 (2) cell lysate.



Western blot analysis using WT1 mouse mAb against K562 (1), COS7 (2), SK-OV-3 (3), HeLa (4), and PC-3 (5) cell lysate.



Flow cytometric analysis of HeLa cells using WT1 mouse mAb (green) and negative control (red).

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com