

DFFB Monoclonal Antibody

Description

Product type	Antibody
Code	BT-MCA3705
Host	Mouse
Isotype	Mouse IgG1
Size	100µL, 50µL
Immunogen	Purified recombinant fragment of human DFFB (AA: 1-289) expressed in E. Coli.
Mol wt	39.1kDa
Species reactivity	Others
Clonality	Monoclonal
Recommended application	IHC,FCM
Concentration	N/A
Full name	N/A
Synonyms	CAD;CPAN;DFF2;DFF40;DFF-40

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene but the biological validity of some of these variants has not been determined.

Recommended Dilution

WB: 1:500 - 1:2000

IHC-p: 1:200 - 1:1000

FCM: 1:200 - 1:400

ELISA: 1:10000

Not yet tested in other applications.

Images

No images.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.