

## ATP1A1 Monoclonal Antibody

### Description

|                                |   |
|--------------------------------|---|
| <b>Product type</b>            | Antibody  |
| <b>Code</b>                    | BT-MCA3763  |
| <b>Host</b>                    | Mouse   |
| <b>Isotype</b>                 | Mouse IgG2a   |
| <b>Size</b>                    | 100μL, 50μL   |
| <b>Immunogen</b>               | Purified recombinant fragment of human ATP1A1 (AA: 153-288) expressed in E. Coli. |
| <b>Mol wt</b>                  | 112.8kDa  |
| <b>Species reactivity</b>      | Others  |
| <b>Clonality</b>               | Monoclonal  |
| <b>Recommended application</b> | WB,IHC,ICC,FCM  |
| <b>Concentration</b>           | N/A   |
| <b>Full name</b>               | N/A   |
| <b>Synonyms</b>                | CMT2DD;HOMGSMR2   |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

### Background

The protein encoded by this gene belongs to the family of P-type cation transport ATPases, and to the subfamily of Na<sup>+</sup>/K<sup>+</sup> -ATPases. Na<sup>+</sup>/K<sup>+</sup> -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The catalytic subunit of Na<sup>+</sup>/K<sup>+</sup> -ATPase is encoded by multiple genes. This gene encodes an alpha 1 subunit. Multiple transcript variants encoding different isoforms have been found for this gene.

### Recommended Dilution

WB: 1:500 - 1:2000

IHC-p: 1:200 - 1:1000


ICC: 1:200 - 1:1000

FCM: 1:200 - 1:400


ELISA: 1:10000

Not yet tested in other applications.


### Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using ATP1A1 mAb against human ATP1A1 (AA: 153-288) recombinant protein. (Expected MW is 40.5 kDa)




Western blot analysis using ATP1A1 mAb against HEK293-6e (1) and ATP1A1 (AA: 153-288)-hlgGFc transfected HEK293-6e (2) cell lysate.




Western blot analysis using ATP1A1 mouse mAb against HeLa (1) and A431 (2) cell lysate.




Immunofluorescence analysis of HeLa cells using ATP1A1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin. Secondary antibody from Fisher (Cat#: 35503)



Flow cytometric analysis of HeLa cells using ATP1A1 mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded esophageal cancer tissues using ATP1A1 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using ATP1A1 mouse mAb with DAB staining.

## Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: [save@bt-laboratory.com](mailto:save@bt-laboratory.com) | [www.bt-laboratory.com](http://www.bt-laboratory.com)