

CD49C Monoclonal Antibody

Description

Product type Antibody

BT-MCA3837 Code

Host Mouse

Mouse IgG1 Isotype $100\mu L$, $50\mu L$ Size

Purified recombinant fragment of human CD49C (AA: extra 63-248) expressed in E. Coli. Immunogen

Mol wt 116.6kDa

Species reactivity Human

Monoclonal Clonality

Recommended application **FCM**

Concentration N/A Full name

N/A

ITGA3;VL3A;FRP-2;GAPB3;ILNEB;MSK18;VCA-2;VLA3a;GAP-B3 Synonyms

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


The gene encodes a member of the integrin alpha chain family of proteins. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain that function as cell surface adhesion molecules. The encoded preproprotein is proteolytically processed to generate light and heavy chains that comprise the alpha 3 subunit. This subunit joins with a beta 1 subunit to form an integrin that interacts with extracellular matrix proteins including members of the laminin family. Expression of this gene may be correlated with breast cancer metastasis.

Recommended Dilution

WB: 1:500 - 1:2000 FCM: 1:200 - 1:400 ELISA: 1:10000


Not yet tested in other applications.

Images




Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)

Western blot analysis using CD49C mAb against human CD49C (AA: extra 63-248) recombinant protein. (Expected MW is 47 kDa)



Western blot analysis using CD49C mAb against HEK293 (1) and CD49C (AA: extra 63-248)-hIgGFc transfected HEK293 (2) cell lysate.



Flow cytometric analysis of HL-60 cells using CD49C mouse mAb (green) and negative control (red).

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com