

4E-BP1 Monoclonal Antibody

Description

Product type Antibody

Code BT-MCA3953

Host Mouse

 Isotype
 Mouse IgG1

 Size
 100µL, 50µL

Immunogen Purified recombinant fragment of 4EBP1 expressed in E. Coli.

Mol wt N/A

Species reactivity Human

Clonality Monoclonal

Recommended application Others

Concentration N/A

Full name N/A

Synonyms BP-1;4EBP1;4E-BP1;PHAS-I;MGC4316;EIF4EBP1

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


4E-BP1(eukaryotic translation Initiation Factor 4E Binding Protein 1),also called ELF4EBP1/BP-1/PHAS-I, which is located on chromosome 8p12, with 118-amino acid protein (about 13kDa). Binding of eIF4EBP1 to eIF4E is reversible and is dependent on the phosphorylation status of eIF4EBP1. Non phosphorylated eIF4EBP1 will bind strongly to eIF4E while(24kDa), the phosphorylated form will not. Akt, TOR, MAP kinase, S6 kinase, and Cdc2 are known kinases capable of inactivating eIF4EBP1 binding to eIF4E by phosphorylating either threonines 35, 45, 69 or serine 64. Although, not all phosphorylation events equally block the eIF4EBP1-eIF4E interaction.

Recommended Dilution

WB: 1:500 - 1:2000 ELISA: 1:10000

Not yet tested in other applications.

Images



Western blot analysis using 4E-BP1 mouse mAb against truncated 4E-BP1 recombinant protein (1) and A431 cell lysate (2).

Storage

Store at $4^{\circ}\mathrm{C}$ short term. Aliquot and store at -20°C long term.