

CD314 Monoclonal Antibody

Description

Product type	Antibody
Code	BT-MCA4063
Host	Mouse
Isotype	Mouse IgG2a
Size	100µL, 50µL
Immunogen	Purified recombinant fragment of human CD314 (AA: extra 73-216) expressed in E. Coli.
Mol wt	25.3kDa
Species reactivity	Others
Clonality	Monoclonal
Recommended application	WB,FCM
Concentration	N/A
Full name	N/A
Synonyms	KLRK1;KLR;NKG2D;NKG2-D;D12S2489E

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Read-through transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster.

Recommended Dilution


WB: 1:500 - 1:2000

FCM: 1:200-1:400


ELISA: 1:10000

Not yet tested in other applications.


Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Western blot analysis using CD314 mAb against human CD314 (AA: extra 73-216) recombinant protein. (Expected MW is 42.6 kDa)




Western blot analysis using CD314 mAb against HEK293-6e (1) and CD314 (AA: extra 73-216)-hIgGFc transfected HEK293-6e (2) cell lysate.




Western blot analysis using CD314 mouse mAb against Rat Spleen (1), A549 (2), and HepG2 (3) cell lysate.



Flow cytometric analysis of Jurkat cells using CD314 mouse mAb (green) and negative control (red).



Flow cytometric analysis of THP-1 cells using CD314 mouse mAb (green) and negative control (red).



Flow cytometric analysis of U937 cells using CD314 mouse mAb (green) and negative control (red).

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com