

NDC80 Monoclonal Antibody

Description

Product type	Antibody
Code	BT-MCA4202
Host	Mouse
Isotype	Mouse IgG1
Size	100μL, 50μL
Immunogen	Purified recombinant fragment of human NDC80 (AA: 443-642) expressed in mammalian.
Mol wt	74KDa
Species reactivity	Human
Clonality	Monoclonal
Recommended application	WB,ICC,FCM
Concentration	N/A
Full name	N/A
Synonyms	HEC;HEC1;TID3;KNTC2;HsHec1;hsNDC80

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

This gene encodes a component of the NDC80 kinetochore complex. The encoded protein consists of an N-terminal microtubule binding domain and a C-terminal coiled-coiled domain that interacts with other components of the complex. This protein functions to organize and stabilize microtubule-kinetochore interactions and is required for proper chromosome segregation.

Recommended Dilution


WB: 1:500 - 1:2000 ICC: 1:200 - 1:1000 FCM: 1:200 - 1:400 ELISA: 1:10000 Not yet tested in other applications.

Images



Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)

Western blot analysis using NDC80 mAb against human NDC80 (AA: 443-642) recombinant protein. (Expected MW is 26 kDa)



Western blot analysis using NDC80 mouse mAb against Hela (1), HepG2 (2), Jurkat (3), K562 (4) and A431 (5) cell lysate.

Immunofluorescence analysis of Hela cells using NDC80 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher (Cat#: 35503)

Flow cytometric analysis of Hela cells using NDC80 mouse mAb (green) and negative control (red).

Storage Store at 4°C short term. Aliquot and store at -20°C long term.

> 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com