

EPCAM Monoclonal Antibody

Description

Product type Antibody

Code BT-MCA4288

Host Mouse

 Isotype
 Mouse IgG1

 Size
 100μL, 50μL

Immunogen Purified recombinant fragment of human EPCAM (AA: extra(116-265)) expressed in E. Coli.

Mol wt 35kDa

Species reactivity Others

Clonality Monoclonal

Recommended application WB,IHC,FCM

Concentration N/A
Full name N/A

 $\textbf{Synonyms} \hspace{35mm} \textbf{ESA;KSA;M4S1;MK-1;DIAR5;EGP-2;EGP40;KS1/4;MIC18;TROP1;EGP314;HNPCC8;TACSTD1} \\$

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy. [provided by RefSeq, Dec 2008]

Recommended Dilution

WB: 1:500 - 1:2000 IHC-p: 1:200-1:1000 FCM: 1:200-1:400 ELISA: 1:10000

Not yet tested in other applications.


Images




Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)

kDa 1 170-130-95-72-55-43-26-17-


Western blot analysis using EPCAM mAb against human EPCAM (AA: extra(116-265)) recombinant protein. (Expected MW is 38 kDa)




Western blot analysis using EPCAM mAb against HEK293-6e (1) and EPCAM (AA: extra(116-265))-hIgGFc transfected HEK293-6e (2) cell lysate.




Western blot analysis using EPCAM mouse mAb against HCT116 (1), HT-29 (2), SW480 (3), Sw-620 (4) , and T47D (5) cell lysate.



Flow cytometric analysis of Lovo cells using EPCAM mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded lung cancer tissues using EPCAM mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded stomach cancer tissues using EPCAM mouse mAb with DAB staining.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com