

HSPA5 Monoclonal Antibody

Description

Product type	Antibody
Code	BT-MCA4309
Host	Mouse
Isotype	Mouse IgG1
Size	100µL, 50µL
Immunogen	Purified recombinant fragment of human HSPA5 expressed in E. Coli.
Mol wt	78kDa
Species reactivity	Human
Clonality	Monoclonal
Recommended application	WB,IHC
Concentration	N/A
Full name	N/A
Synonyms	BIP;MIF2;GRP78;FLJ26106;HSPA5

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

When Chinese hamster K12 cells are starved of glucose, the synthesis of several proteins, called glucose-regulated proteins (GRPs), is markedly increased. Hendershot et al. (1994) (PubMed 8020977) pointed out that one of these, GRP78 (HSPA5), also referred to as 'immunoglobulin heavy chain-binding protein' (BiP), is a member of the heat-shock protein-70 (HSP70) family and is involved in the folding and assembly of proteins in the endoplasmic reticulum (ER). Because so many ER proteins interact transiently with GRP78, it may play a key role in monitoring protein transport through the cell. Probably plays a role in facilitating the assembly of multimeric protein complexes inside the ER. The HSP70 proteins are ubiquitous molecular chaperones that are found in all organisms and tissue types. Like other members of the HSP70 family, BiP is a peptide-binding ATPase that is able to differentiate native proteins from unfolded polypeptides. BiP does not bind to fully folded and assembled proteins, except in the presence of other co-chaperones. BiP is involved in a number of key mechanisms and pathways including polypeptide translocation across the endoplasmic reticulum, folding, assembly, transport of secreted or membrane proteins, and the regulation of calcium homeostasis. Although BiP is relatively abundant, marked increases in BiP occur where there is an accumulation of unfolded polypeptides. For this reason, BiP has been identified as a marker for various disease states that are associated with secretory and transmembrane protein misfolding.

Recommended Dilution


WB: 1:500 - 1:2000

IHC-p: 1:200 - 1:1000


ELISA: 1:10000

Not yet tested in other applications.

Images



Western blot analysis using HSPA5 mouse mAb against NIH/3T3 (1), HeLa (2) and Jurkat (3) cell lysate.



Immunohistochemical analysis of paraffin-embedded human Thyroid tissues using HSPA5 mouse mAb

Storage

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com