

MLH1 Monoclonal Antibody

Description

Product type	Antibody
Code	BT-MCA4319
Host	Mouse
Isotype	Mouse IgG1
Size	100μL, 50μL
Immunogen	Purified recombinant fragment of human MLH1 (AA:381-483) expressed in E. Coli.
Mol wt	84.6kDa
Species reactivity	Others
Clonality	Monoclonal
Recommended application	WB,IHC,FCM
Concentration	N/A
Full name	N/A
Synonyms	FCC2;COCA2;HNPCC;hMLH1;HNPCC2

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

The protein encoded by this gene can heterodimerize with mismatch repair endonuclease PMS2 to form MutL alpha, part of the DNA mismatch repair system. When MutL alpha is bound by MutS beta and some accessory proteins, the PMS2 subunit of MutL alpha introduces a single-strand break near DNA mismatches, providing an entry point for exonuclease degradation. The encoded protein is also involved in DNA damage signaling and can heterodimerize with DNA mismatch repair protein MLH3 to form MutL gamma, which is involved in meiosis. This gene was identified as a locus frequently mutated in hereditary nonpolyposis colon cancer (HNPCC). [provided by RefSeq, Aug 2017]

Recommended Dilution


WB: 1:500 - 1:2000 IHC-p: 1:200-1:1000 FCM: 1:200-1:400 ELISA: 1:10000 Not yet tested in other applications.

Images



Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)

Western blot analysis using MLH1 mAb against human MLH1 (AA: 381-483) recombinant protein. (Expected MW is 25.1 kDa)



Western blot analysis using MLH1 mAb against HEK293 (1) and MLH1 (AA: 381-483)-hIgGFc transfected HEK293 (2) cell lysate.

Store at 4°C short term. Aliquot and store at -20°C long term.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com